

SYMPOSIUM

Free radical properties of metal complexes*

Klára Szentmihályi^{1*}, Anna Blázovics^{1,2}, Péter Vinkler¹

¹Chemical Institute, Chemical Research Center, Hungarian Academy of Sciences, Budapest, Hungary, ²2nd Department of Medicine, Semmelweis University, Budapest, Hungary

ABSTRACT Since metals are capable of catalysing the free radical reactions, different metal (Mg, Mn, Zn) compounds were studied to determine their hydroxyl radical generating and/or scavenging abilities depending on ligand in a H_2O_2/OH^- -luminol system with or without microperoxidase. Magnesium oxide, magnesium gluconate and manganese gluconate increase the free radical reactions in the H_2O_2/OH^- -luminol system. Most of examined metal compounds have different hydroxyl radical scavenging activity in H_2O_2/OH^- -microperoxidase-luminol system. Magnesium citrate has the highest antioxidant effect, while manganese compounds seem to be prooxidant. Vitamin E and C elevate the free radical level in this experimental system.

Acta Biol Szeged 47(1-4):107-109 (2003)

KEY WORDS

chemiluminescence
free radical generating ability
scavenging activity
magnesium, manganese and zinc
compounds

In living organisms free radical formation and the free radical defense system are closely related to metal ions and complexes (Tampo and Yonaha 1992; Sigel and Sigel 1999). Fenton reaction describes Fe(II) induced hydroxyl radical formation (Fenton 1894, Haber and Willstätter 1931, Haber and Weiss 1934) and other transition metal ions (e.g. V, Cr) take part in Fenton-like reactions (Halliwell and Gutteridge 1990; Goldstein et al. 1993). Nevertheless, none transition metal ions also play a significant role, directly or indirectly, in free radical reactions. For example manganese, zinc and copper have a role in the antioxidant defense system (Zidenberger-Cherr and Keen 1991; Ebadi et al. 1996) and magnesium is present in every cell membrane in membrane ATP-ase and magnesium activates several enzymes which catalyse polyphosphate hydrolysis (Fox et al. 2001).

In the diseases where metal ion supplementation is necessary, it is important to know the complex form (ligand) of the metal. The absorption of metal and other physico-chemical properties can change, depending on the ligand (Hallberg et al. 1970; Kaltwasser et al. 1987; Rauch et al. 1990; Szentmihályi et al. 1998) and the physiological effect of the metal compound varies depending on the ligand as well.

Therefore, the hydroxyl radical generating ability and scavenging activity of some metal (Mg, Mn, Zn) compounds were studied by chemiluminescence method in the H_2O_2 -luminol system with or without microperoxidase.

Materials and Methods

The weight ratio of vitamin E to vitamin C was 1:2, and that of vitamins to metals was 2:1 in the experimental system. Concentrations of the metal compounds in the basic solutions were 1 mg/ml. The compounds examined were of pharmacopoeal quality. Since the result is matrix dependent, the measurements were always made under same conditions (temperature, pH, concentration ratios).

Chemiluminescence techniques were applied for the studying of metal catalysis or scavenger activity. Light emission of luminol was measured by a method of Blázovics et al. (1999) using a Berthold Lumat LB-9501 luminometer. The intensity of chemiluminescence light is given as the relative light unit (RLU).

The reaction mixture for measuring free radical generating ability was as follows: hydrogen peroxide (10^4 dilution) 300 μ l, luminol (7×10^{-7} M) 100 μ l, the sample (1 mg metal compound/ml solution) 50 μ l and bidistilled water 250 μ l.

The reaction mixture for measuring the hydroxyl scavenging activity was: hydrogen peroxide (10^4 dilution) 300 μ l, microperoxidase (3×10^{-7} M) 300 μ l, luminol (7×10^{-7} M) 50 μ l and the sample (1 mg/ml) 100 μ l diluted with bidistilled water to 1 ml. The intensity of chemiluminescence light is given as the relative light unit (RLU) reduced by free radical scavenging substances.

Results and Discussion

Since the decomposition of luminol in the presence of hydrogen peroxide is catalysed by metal ions, some metal compounds were investigated for the catalytic activity of the reaction.

Accepted April 30, 2003

*Corresponding author. E-mail: szklari@chemres.hu

[†]In memory of Profesor Béla Matkovic

Table 1. Free radical generating or scavenging abilities of metal compounds (RLU %) and standard deviation (SD %) in the H₂O₂/•OH- luminol system.

Samples	RLU %	SD %
Background int	100	
Magnesium oxi	125	1.2
Magnesium chl	71.0	0.8
Magnesium sul	20.6	0.4
Magnesium cit	5.1	1.0
Magnesium glu	125	1.1
Manganese dic	26.8	1.3
Manganese sul	12.5	0.3
Manganese glu	102	1.5
Zinc chloride	4.9	0.6
Zinc sulfate	6.6	0.2
Zinc citrate	13.5	1.7
Zinc gluconate	6.0	1.4
Vitamin E and C	15.6	0.5

Table 2. Free radical generating or scavenging abilities of metal compounds (RLU %) and standard deviation (SD %) in the H₂O₂/•OH- microperoxidase-luminol system.

Samples	RLU %	SD %
Background int	100	
Magnesium oxi	177.4	0.9
Magnesium chl	104.7	0.9
Magnesium sul	106	0.3
Magnesium cit	2.3	2.5
Magnesium glu	388	4.9
Manganese dic	364	2.5
Manganese sul	366	0.4
Manganese glu	203	3.0
Zinc chloride	201.6	1.1
Zinc sulfate	82.6	0.1
Zinc citrate	74.3	3.6
Zinc gluconate	82.1	6.6
Vitamin E and	91.6	0.4

Table 3. Free radical generating or scavenging abilities of metal compounds in the presence of Vitamin E and C (RLU %) and standard deviation (SD %) in the H₂O₂/•OH- microperoxidase-luminol system

Samples	RLU %	SD %
Background intensity	100	
Magnesium sulfate	191.5	1.2
Magnesium gluconate	182.9	0.8
Manganese sulfate	240	0.3
Manganese gluconate	160.5	0.4
Zinc sulfate	95.9	0.1
Zinc gluconate	94.4	0.4

In the applied system and concentrations neither vitamins nor metal compounds (except for magnesium oxide, magnesium gluconate and manganese gluconate) generated the decomposition of luminol to aminophthalate in H₂O₂/•OH- luminol system (Table 1). When we applied the microperoxidase in the system (H₂O₂/•OH-microperoxidase-luminol) different metal compounds have different radical scavenging activity (Table 2). The metal ions, depending on the quality of the metal, ligand and concentration, may induce free radicals. Among magnesium compounds, only magnesium citrate has significant scavenging activity whereas other magnesium compounds enhance the free radical level formation. None of the manganese compounds have hydroxyl radical scavenging activity, while zinc compounds (except for zinc chloride) show scavenging activity. The mixture of vitamin E and C in the applied concentration has only a weak scavenging activity in this system.

The presence of vitamin E and C in the metal ion-containing solution changes the free radical scavenging activity. The vitamins increased the relative light unit (RLU %) of the system (H₂O₂/•OH-microperoxidase-luminol) together with magnesium sulfate, zinc sulfate and zinc gluconate solutions, while decreased free radical formation together with magnesium gluconate, manganese sulfate and manganese gluconate solutions (Table 3).

According to the results it has been stated that chemiluminescence measurement is an efficient method for the determination of free radical generating or scavenging activity of metal compounds as well.

Acknowledgments

This work was supported by the National Research and Development Program 1/047/2001.

References

- Blázovics A, Kovács Á, Lugasi A, Hagymási K, Bíró L, Fehér J (1999) Antioxidant defence in erythrocytes and plasma of patients with active and quiescent Crohn's disease and ulcerative colitis: A chemiluminescent study. *Clin Chem* 45:895-896.
- Ebadi M, Leuschen MP, Refaey H, Hamada F M, Rojas P (1996) The antioxidant properties of zinc and metallothionein. *Neurochem Int* 29:159-166.
- Fenton HJH (1894) Oxidation of tartaric acid in presence of iron. *J Chem Soc* 65:899-910.
- Fox C, Ramsoomair D, Carter C (2001) Magnesium: its proven and potential clinical significance. *Southern Med J* 94:1195-1201.
- Goldstein S, Meyerstein D, Czapski G (1993) The Fenton reagents. *Free Rad Biol Med* 15:435-445.
- Haber F, Willstätter R (1931) Unpairedness and radical chains in the reaction mechanism of organic and enzymic process. *Ber Deutsch Chem Ges* 64:2844-2856.
- Haber F, Weiss J (1934) The catalytic decomposition of hydrogen peroxide by iron salts. *Proc Roy Soc London*, 147:332-351.
- Hallberg L, Harwerth HG, Vanotti A eds., (1970) Iron deficiency: pathogenesis, clinical aspects, therapy. Academic Press, London, New York, 1-628.

- Halliwell B, Gutteridge JMC (1990) Role of free radicals and catalytic metal ions in human disease: an overview. *Meth Enzymol* 86:1-85.
- Kaltwasser JP, Werner E, Niechzial M (1987) Bioavailability and therapeutic efficacy of bivalent and trivalent iron preparations. *Arz-Forsch Drug Res* 37:122-129.
- Rauch P, Kas J, Ranny M (1990) iron bioavailability from its complex with sucrose. *Food Chem* 36:129-134.
- Sigel A, Sigel H (1999) Interactions Between Free Radicals and Metal Ions in Life Process. In Sigel A, Sigel H eds., *Metal Ions in Biological Systems*. Marcel Dekker Inc, New York, Basel, 36:1-797.
- Szentmihályi K, Lakatos B, Sándor Z, Hajdú M, Vinkler P (1998) Preparation of magnesium polygalacturonate complexes for magnesium supplementation and their physicochemical characterization. In Kiss AS ed., *Magnesium. Magnesium and Interaction of Magnesium with Trace Elements*. Hungarian Chemical Society, Budapest, 241-244.
- Tampo Y, Yonaha M (1992) Antioxidant mechanism of manganese (II) in phospholipid peroxidation. *Free Radical Biol Med* 13:115-120.
- Zidenberger-Cherr S, Keen CL (1991) Essential trace elements in antioxidant processes. In Dreosti IE ed., *Trace Elements, Micro Nutrients, and Free Radicals*. Humana Press, Totowa, New Jersey, 107-127.