

Light response of the chlorophyll fluorescence parameters and partitioning of absorbed light energy in wild type and *npq4* mutant of *Arabidopsis thaliana*

Zsuzsanna Várkonyi^{1*}, Szilvia Bajkán, Gyula Váradi², Endre Lehoczki

Department of Botany and Botanical Garden, University of Szeged, Szeged, Hungary, ¹Institute of Plant Biology, Biological Research Centre, Hungarian Academy of Sciences, Szeged, Hungary, ²Research Institute for Viticulture and Enology, Kecskemét, Hungary

ABSTRACT Photoprotective thermal dissipation of absorbed light energy is measured as non-photochemical quenching (NPQ) of chlorophyll fluorescence during exposure of plants to higher light intensities. We investigated the light response behavior of chlorophyll fluorescence and energy partitioning in PSII of *npq4* mutant of *Arabidopsis thaliana*, which is lacking of PsbS protein, to that of the wild type plants. We found that the PsbS protein appears to function not only in thermal dissipation in light via xanthophyll pigments, but also modified the allocation pattern of excitation energy absorbed in the PSII to PSII photochemistry (P), thermal dissipation in light (DL) and excess excitation (E).

Acta Biol Szeged 49(1-2):229-232 (2005)

KEY WORDS

Arabidopsis
npq4 mutant
chlorophyll fluorescence
energy allocation

Photosynthetic organisms are exposed to a wide range of light intensities in nature. Above a certain level photosynthesis is incapable of using all of the energy absorbed by the light-harvesting complexes (LHCs). To prevent the damage caused by excess excitation various photoprotective mechanisms are present in the thylakoid membrane. One of these mechanisms is thermal dissipation of the excess excitation energy within the LHCII (Demmig-Adams et al. 1996; Elrad et al. 2002)

Photoprotective thermal dissipation of absorbed light energy is measured as non-photochemical quenching (NPQ) of chlorophyll fluorescence during exposure of plants to higher light intensities. In wild-type *Arabidopsis* and other plants several processes can contribute to NPQ; the major component is Δ pH-dependent NPQ, called qE. Following the induction of qE by light, the alteration of Δ pH activates synthesis of zeaxanthin, through the xanthophyll cycle. The conformational changes in the structure of LHCII, especially in PsbS protein caused by binding of protons and zeaxanthin to PsbS induce quenching of excitation energy. The *NPQ4* gene encodes the PsbS protein what is a member of the chlorophyll a/b-binding, light harvesting complex family of proteins and binds zeaxanthin. Genetic analysis showed that the phenotype of *npq4-1* mutant plants was due to a single, nuclear mutation, which blocks qE that is detected in the wild type plants. The *npq4-1* mutant lacks the Δ pH- and zeaxanthin-dependent conformational change in the thylakoid membrane that is necessary for qE (Li et al. 2002).

In the present study, we compared the light response behavior of chlorophyll fluorescence and energy partitioning in PSII of *npq4* mutant of *Arabidopsis thaliana* to that of the wild type plants.

*Corresponding author. E-mail: photosynt@nucleus.szbk.u-szeged.hu

Materials and Methods

Plant material

Seeds were germinated and grown in commercial potting soil mixture in the growth chamber, with 12 h /12 h day/night regime, with corresponding temperature 25°C/22°C under an irradiance of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photon flux density (PFD).

Chlorophyll fluorescence measurements

Chlorophyll fluorescence measurements were carried out in dark adapted leaves with a pulse amplitude modulated fluorometer (PAM 200 Walz, Germany). In the measurements the standard setting of Run 8 were used according to the PAM-200 handbook (Operating Manual, 1st Edition, 1997). Before start of the Run 8, initial fluorescence yield (F_0) and maximum fluorescence yield (F_m) were determined in 60 min dark adapted samples with a saturating pulse (3000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PFD). After start of the Run 8, a 5 min preillumination at 210 $\mu\text{mol m}^{-2} \text{s}^{-1}$ actinic light (AL) intensity was applied in order to allow light adaptation of the sample and activation of Calvin cycle enzymes. Then AL changed to 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PFD and increased every 5 min until 1250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PFD has reached (10 steps). The quenched levels of maximum chlorophyll fluorescence (F_m') parameters were determined at each AL level by saturated pulses (3000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PFD, of 1 s duration) applied to the leaves at the end of 10 s AL illumination. After the AL have been switched off, far-red light was exerted for the determination of the minimal (F_0') level of fluorescence for correct determination of qP and F_v'/F_m' at each light intensity. The terminology suggested by van Kooten and Snel (1990) was used. From measured parameters (F_0 , F_m , F_0' , F_m' and the steady state fluorescence level, F_s) the




Figure 1. Actinic light response of the photochemical quenching (A), the yield of PSII photochemistry (B), the quantum yield of open PSII centers (C) and the non-photochemical quenching (D) in leaves of wild and *npq4* mutant of *A. thaliana*. The data are means \pm SE of minimum 6 replicates from 3 independent experiments.




Figure 2. Light response allocation of light absorbed by the PSII antenna to PSII photosynthetic electron transport (area of P), thermal energy dissipation in the light (area of DL) and in the dark (area of DD), and to excess excitation (area of E) in the leaves of wild (A) and *npq4* mutant (B) of *A. thaliana*. The data are calculated from the average values of the results presented in Figure 1.

Table 1. Pigment composition, sizes of the xanthophyll cycle pool and de-epoxidation index (DEi) of the leaves of wild and *npq4* mutant type of *Arabidopsis thaliana*. The results are the means \pm SE of the three independent experiments.

	Chl-a+b ($\mu\text{mol m}^{-2}$)	Caroten. ($\mu\text{mol m}^{-2}$)	Chl-a/b	Chl-a+b/ Carotenoids	V+A+Z ($\mu\text{mol m}^{-2}$)	V+A+Z/Chl-a+Chl-b ($\mu\text{mol mmol}^{-1}$)	DEi
Wild	129 \pm 4	41 \pm 4	2,57 \pm 0,1	3,2 \pm 0,1	12 \pm 1,6	93 \pm 6	0,60 \pm 0,03
Mutant	128 \pm 3	46 \pm 5	2,87 \pm 0,2	2,8 \pm 0,2	12,5 \pm 1,1	98 \pm 4	0,58 \pm 0,03

optimum quantum yield (Fv/Fm), photochemical quenching coefficient (qP), non-photochemical quenching (NPQ) and relative quantum yield of PSII photochemistry (QY) were calculated according to Schreiber et al. 1986; Genty et al. 1989; Bilger and Björkman 1990, respectively. The partitioning of absorbed light energy in PSII was calculated according to the model of Demmig-Adams et al. 1996)

Pigment analysis

Chlorophylls and total carotenoids were measured according to Lichtenthaler (1987). Xanthophyll cycle components were determined by means of HPLC (Váradi et al. 1992).

Results and Discussion

In dark-adapted leaves the optimum quantum yield of PSII was $0,83 \pm 0,02$ for both types of *A. thaliana*. Figure 1 shows the light response of calculated chlorophyll fluorescence parameters. The fraction of open PSII centers (qP, Fig. 1A), the quantum yield of PSII electron transport (QY, Fig. 1B) and the quantum yield of open PSII (Fv'/Fm', Fig. 1C) decreased with increasing PFD, but more sensitively with the light intensity in the *npq4* mutant plants. Only values of Fv'/Fm' at higher PFDs in mutant plant were slightly higher. Moreover, as expected, significantly lower level of NPQ was observed in *npq4* mutant of *A. thaliana* (Fig. 1D). However, under the growth conditions employed, pigment composition and the *in vivo* levels of the xanthophyll pigments violaxanthin (V), antheraxanthin (A) and zeaxanthin (Z) in excess light were the same for *npq4* mutant and wild type of *A. thaliana*, and showed very similar xanthophyll de-epoxidation (DEi) activity (Table 1).

For a better understanding the possible role of PsbS protein in the allocation of absorbed light energy to the PSII photochemistry and thermal dissipative processes was assessed for the wild and *npq4* mutant of *A. thaliana* with the model of Demmig-Adams et al. (1996). The fraction of light absorbed by PSII antennae that is utilized in the PSII photochemistry was estimated from $P=(Fv'/Fm') qP$. Heat dissipation is defined as $1-Fv'/Fm'$, which was divided in to the fraction of energy dissipated in the dark ($DD=(1-Fv'/Fm')$) and that dissipated in the light ($DL=Fv'/Fm'-Fv'/Fm'$) via xanthophyll cycle. The fraction of absorbed light neither going into P, nor into DD and DL is defined as excess (E), and it is calculated as the product of (Fv'/Fm') ($1-qP$). The

calculated energy partitioning is shown in Fig. 2. In both type of plants the fraction of photochemistry (P) decreased, but DL and E increased with increasing PFD. The *npq4* mutant showed slightly lower DL, significantly lower P, and considerably higher E values. It seems, that the decrease in DL and P contributed to the increase in excess energy, which is also a consequence of PsbS protein mutation.

The PsbS protein is necessary for photoprotective thermal dissipation of excess absorbed light energy in plants, measured as non-photochemical quenching of chlorophyll fluorescence. However, the ΔpH and xanthophyll cycle-dependent conformational change associated with qE is absent in *npq4* mutant. It is well-known that *npq4* mutant of *A. thaliana* that lack qE is more sensitive to light stress. Our results suggest that the PsbS protein appears to function not only in thermal dissipation in light via xanthophyll pigments, but also modified the allocation pattern of excitation energy absorbed in the PSII to PSII photochemistry (P), thermal dissipation in light (DL) and excess excitation (E).

Acknowledgments

We thank for the *npq4* mutant of *Arabidopsis thaliana* seeds for prof. P. Horton, Sheffield.

This work was supported by OTKA grant T035252, which is gratefully acknowledged.

References

- Bilger W, Björkman O (1990) Role of the xanthophyll cycle in photoprotection elucidated by measurements of light induced absorbance, fluorescence and photosynthesis in leaves of *Hedera canariensis*. *Photosynth Res* 25:173-185.
- Demmig-Adams B, Adams III WW, Barker DH, Logan BA, Bowling DR, Verhoeven AS (1996) Using chlorophyll fluorescence to assess the fraction of absorbed light allocated to thermal dissipation of excess excitation. *Physiol Plantarum* 98:253-264.
- Elrad D, Niyogi KK, Grossman AR (2002) A major light-harvesting polypeptide of photosystem II functions in thermal dissipation. *The Plant Cell* 14:1801-1816.
- Genty B, Briantais JM, Baker NR (1989) The relationship between the quantum yield of non-photochemical quenching of chlorophyll fluorescence and the rate of photosystem 2 photochemistry in leaves. *Biochim Biophys Acta* 990:87-92.
- Li XP, Björkman O, Shih C, Grossman AR, Rosenquist M, Jansson S, Niyogi KK (2000) A pigment-binding protein essential for regulation of photosynthetic light harvesting. *Nature* 403:391-395.
- Li XP, Phippard A, Pasari J, Niyogi KK (2002) Structure-function analysis of photosystem II subunit S (PsbS) *in vivo*. *Funct Plant Biol* 29:1131-1139.

- Lichtenthaler HK (1987) Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods Enzymol* 148:350-382.
- Schreiber U, Schliwa U, Bilger W (1986) Continuous recording of photochemical and non-photochemical quenching of chlorophyll fluorescence with a new type of modulation fluorometer. *Photosynth Res* 10:51-62.
- van Kooten O, Snel JFH (1990) The use of fluorescence nomenclature in plant stress physiology. *Photosynth Res* 25:147-150.
- Váradi Gy, Botos-Bálo B, Pölös E (1992) Xantophyll cycle in grapevine leaves – diurnal and seasonal patterns. *Proceedings of the IVth International Symp on Grapevine Physiology*, 11-15 May. San Michelle All'adige, Torino 521-526.