

ARTICLE

Antioxidant, physicochemical and mineral evaluations of *Spondias mombin* crude fruit juice

Akharaiyi Fred Coolborn^{1*}, Boboye Bolatito Esther², Akinyosoye Felix Akinsola²
Oso Benjamin Afolabi¹

¹Department of Biological Sciences, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria

²Department of Microbiology, The Federal University of Technology, Akure, Ondo State, Nigeria

ABSTRACT Edible fruits have potential health benefits regarding their richness in minerals, nutrients, vitamins and antioxidant. In view of the medicinal potential of the leaves and stem bark extracts of *Spondias mombin*, this study aimed to determine the antioxidant, physicochemical and mineral contents of the crude fruit juice of *S. mombin* for its employment in health promotion and traditional use for medicine. Free radical scavenging power of 93.97 ± 64.8 μmol (TE) and Ferric reducing power (FRAP) of 11.8 ± 0.2 μmol (AAE) were observed *in vitro* from the crude fruit juice. The physicochemical property of the crude fruit juice yielded proximate compositions of protein (6.03 ± 0.44), fat (1.85 ± 0.03), pH (3.63 ± 0.11), fibre (0.64 ± 0.02), moisture ($80.60 \pm 0.22\%$), ash (0.54 ± 0.03), carbohydrate (11.61 ± 0.34) and titratable acidity (28.52 ± 0.14). Trace elements from the crude fruit juice are in the amounts of 276.27 ± 2.65 for potassium, 1.15 ± 0.01 (manganese), 0.121 ± 0.32 (copper), 4.45 ± 1.42 (sodium), 136.42 ± 1.35 (zinc), 12.36 ± 0.03 (magnesium), 0.01 ± 0.00 (lead), 28.22 ± 1.02 (phosphorus), 13.04 ± 0.53 (calcium) and 0.48 ± 0.01 (iron).

Acta Biol Szeged 60(2):171-176 (2016)

KEY WORDS

antioxidant
fruit juice
physicochemical
Spondias mombin

Introduction

In Nigeria, large numbers of underutilized wild fruits are widely distributed throughout the country. Most of these wild fruits are consumed mainly by rural dwellers, who appreciate them in quenching of thirst and supplement for food without knowledge about their health importance and medicinal values. The Nigeria climate has favorable potency for the production of tropical fruits. These tropical fruits are seasonal and some are cultivated for their health benefits, while others are readily available as wild in the forests uncultivated by man. Fruits are important sources of many nutrients and have served as integral part of the human diet as they supply vitamins, minerals and other vital constituents essential for human health (Akhtar et al. 2010). Fruits are believed to occupy a modest place as a source of trace elements due to their high water content. However, consumers look for variety in their diets and therefore opt for fresh fruits because of the health benefits. Fruits are special food sources that are rich in potassium (K), calcium (Ca) and magnesium (Mg). These

minerals among others are nutrient requirements of which their intake is associated with reduced risk of, e.g., cardiovascular diseases, cataracts, and age-related functional decline (Willett 1994, 1995; Temple 2000; Aberoumand and Deokule 2010). Higher amounts of antioxidants such as vitamin A, C, E, lycopene, polyphenols and carotene are present in fresh fruits, than when prepared into juice and beverages these components are often reduced.

Numerous studies have shown that fruits are rich sources of nutrients, as well as non-nutrient molecules with antioxidants or other physiological effects that are important constituents of a healthy diet. The health promoting properties of plant-based foods have largely been attributed to their wide range of phytochemicals, of which many are present at relatively high levels (Ali et al. 2011).

Normal physiological processes *in vivo* result in the production of free radicals. Oxidative stress created when there is an insufficient capacity of the biological system to neutralize excess free radicals. This may result in aging and disease conditions as reported by Sahreen et al. (2014). Fruits and other foods are known to contain antioxidants that are linked to *in vivo* protection from oxidative stress (Jensen et al. 2008). Different studies have shown that free radicals present in the human organs, cause oxidative damage to various bio-molecules, such as lipids, proteins and nucleic acids, and

Submitted October 10, 2016; Accepted November 23, 2016

*Corresponding author. E-mail: akharaiyifc@abuad.edu.ng

thus are involved in the initiation phase of certain degenerative diseases. Phenolic and other phytochemical antioxidants found in fruits and vegetables capable of neutralizing free radicals and may play a major role in the prevention of certain diseases (Kaur and Kapoor 2001).

Numerous epidemiological investigations have pointed out that a lack of essential mineral elements in food can result in an increase in sensitivity to illness, leading to suboptimal health or an enhancement of disease occurrence. Among others iodine, selenium, zinc, iron, copper, manganese and chromium were recognized as essential mineral elements indispensable for maintaining normal life activities. All these mineral elements can be obtained from the consumption of certain foods. It was also recognized decades ago that green and natural foods of plant origin are of important sources of these minerals. These metals are commonly found in trace amounts in various plants as they easily take up them from the environment. At the same time, different parts of a plant could contain different concentration of these elements. In spinach, e.g., the amount of Mn in the tip of the leaf was higher than in the other parts, or Co and Zn were mainly accumulated around the main veins (Xin et al. 2009).

Spondias mombin is a tree having habitat in several tropical forests in the world. In Nigeria, it is widely distributed in all geographical regions as wild plant. It is called *Iyeye* by the Yorubas, *Ichikara*, or *Uvuru* by the Ibos, and *Tsaadar lamarudu* by the Hausas. Names of this plant in some other parts of the world are *Bala* (Costa Rica), *Jobito* (Panama), *Jobo blanco* (Colombia), *Jobo corronchoso* (Venezuela), *Hoeboe* (Surinam), *Acaiba*, *Caja*, *Pau da taperra* (Brazil), *Ubo* (Peru), and *Hobo* (Mexico). For its ethno-medicinal uses, the fruit juice is consumed as a diuretic, febrifuge and laxative. The aims of this study were to analyze the antioxidant activity, physicochemical properties and mineral composition of *S. mombin*. The long-term goal is to promote and increase the utilization and consumption of edible wild fruits.

Materials and Methods

Collection and preparation of fruit juice

Ripe fruits of *S. mombin* were harvested from its tree at the town of Ado Ekiti (Nigeria). The fruit was identified by a botanist in the Department of Biological Sciences, Afe Babalola University (Ado Ekiti, Nigeria). The fruits were washed with 3% sodium hypochlorite and rinsed severally in distilled water. The fruits were pressed for juice to be extracted. The extracted crude juice was filtered through No 1 Whatman filter paper and the filtered juice was kept in a sterile bottle at 4°C before use.

Determination of radical scavenging capacity with 1,1-diphenyl-2-picrylhydrazyl (DPPH)

DPPH radical scavenging activity was determined according to Cavin et al. (1998). Two milliliters of the crude fruit juice was added to 1 ml of DPPH solution (0.04 mg/ml in methanol). The mixture was vigorously shaken and incubated in the dark for 20 min. Thereafter, the reduction of DPPH absorption was measured at 517 nm. A calibration curve was prepared by measuring the reduction in absorbance of the DPPH solution in the presence of different concentrations of Trolox (0-400 µM). Results were expressed as µmol of Trolox equivalents (TE)/mg wet extract. All determinations were performed in triplicate.

Determination of ferric reducing activity (FRAP)

The ferric reducing activity of the fruit juice was estimated based on the FRAP assay (Benzie and Strain 1996). The solutions for this assay are 300 mmol/L acetate buffer, 10 mmol/L TPTZ (2,4,6- tris(2-pyridyl)-s-triazine) in 40 mmol/L of HCl and 20 mmol/L FeCl₃ (6 H₂O). From this 25 ml acetate buffer was mixed with 2.5 ml of TPTZ solution and 2.5 ml FeCl₃ (6 H₂O). The assay was performed by adding 1 ml of the FRAP reagent to 2 ml of diluted fruit juice. Absorbance reading at 593 nm was recorded after 20 min of the reaction. The change in absorbance was related to the absorbance change in ascorbic acid as the standard solution tested in parallel. Results were expressed as micromole of ascorbic acid equivalents (AAE)/mg of wet weight of the fruit juice. All determinations were performed in triplicate.

Determination of total phenol

Total phenol concentration of the crude fruit juice was determined by spectrophotometric method, according to the Folin-Ciocalteu colorimetric method of Singleton and Rossi (1965), using caffeic acid as the standard and expressing the results as µmol equivalents of caffeic acid (CAE)/mg of wet weight of fruit juice.

Determination of total flavonoids content

Total flavonoids from the crude fruit juice were estimated by colorimetric method using the criteria of Sakanaka et al. (2005). Aliquot of fruit juice was mixed with 400 µl of distilled water and 30 µl of a 5% sodium nitrite solution. After 6 min, 10% aluminium chloride solution (30 µl) was added and the mixture was allowed to stand for 5 min. Then 200 µl of 1M NaOH solution and 240 µl of distilled water were added and the absorbance was measured at 510 nm. Rutin was used as standard solutions. All determinations were done in

Table 1. Antioxidant profile of the crude fruit juice of *S. mombin*.

Antioxidant	Mean \pm SD
Free radical scavenging activity (DPPH) ($\mu\text{mol TE}$)	93.97 \pm 64.8
Ferric reducing activity (FRAP) ($\mu\text{mol AAE}$)	11.8 \pm 0.24
Total flavonoids (RE/mg wet fruit juice)	14.5 \pm 0.04
Total phenol (CAE/mg wet weight fruit juice)	155.126 \pm 12.546

triplicate and values were calculated from a calibration curve obtained with rutin. Final results were expressed as μmol of rutin equivalents (RE)/mg wet fruit juice.

Determination of physicochemical properties

The standard methods of Association of Official Analytical Chemists, (AOAC 2005) was used to analyze the physicochemical properties of the *S. mombin* crude fruit juice. All analysis were made in triplicate.

Determination of mineral contents

Wet ashing method was used to digest the crude fruit juice and the quantification was by atomic spectroscopy and mass spectrometry (ICP-MS, Spectro Analytical Instruments GmbH, Germany) according to the AOAC methods 997.15 and 990.08 (AOAC 2005). The results were expressed in mg/100 g. All analysis were performed in triplicate.

Statistical analysis

The results were expressed as mean \pm standard deviation (SD) and subjected to one way analysis of variance (ANOVA). The least significant difference (LSD) was performed for the pairwise mean comparisons to determine the significant treatment dose at 95% level of confidence. Values were considered statistically significant at ($P < 0.05$).

Results and Discussion

Antioxidants capacity of the crude fruit juice of *S. mombin*

Free radical scavenging power of $93.97 \pm 64.8 \mu\text{mol TE}$ and ferric reducing power (FRAP) of $11.8 \pm 0.2 \mu\text{mol AAE}$ were measured as reasonable amounts of the juice ability in *in vitro* antioxidant capacity (Table 1). The total antioxidant activity shown by the crude fruit juice of *S. mombin* reflects

substantial free radical scavenging ability and reductive potential.

Free radicals are frequently generated in the living system and could be responsible for cell and tissues damage. As a consequence, research is focused on exploring for safe and effective antioxidants and also to encourage the consumption of natural antioxidants from food supplements and traditional medicines (Yazdanparast and Ardestani 2007). Antioxidant compounds, such as phenolic acids, carotenes and vitamins have been reported to be naturally present in fruits, vegetables, nuts, herbs and spices (Ali et al. 2008; Vasco et al. 2008; Schinella et al. 2009; Sreeramulu and Raghunata 2010; Lu et al. 2011). The results obtained in this study, indicates that the crude fruit juice of *S. mombin* is a good source of antioxidants, that could be of help to alleviate oxidative stress. Based on the evaluated antioxidant properties by Adewale et al. (2015), Famobuwa et al. (2016a; 2016b) on some Nigerian fruits, the result obtained allows us the insight that the crude fruit juice of *S. mombin* possessed antioxidant activity above average. Along with some fruits, such as pineapple, banana, watermelon, and orange and vegetables such as tomato and carrot, that are commonly consumed on daily bases by the populace. Also, the report of free radical scavenging activity (71.5%) and ferric reducing potential (1.94%) from the leaves of *S. mombin* as reported by Awogbindin et al. (2014) were less value to what was obtained from the crude fruit juice of *S. mombin* in this study.

The amounts of phenolic compounds and flavonoids contents of *S. mombin* in this study suggest its high antioxidant capacity. Phenolic compounds of plants fall into several categories; chief among these are the flavonoids, which have potent antioxidant activities (Tadhani et al. 2007). Flavonoids derivatives have shown a wide range of antibacterial, antiviral, anti-inflammatory, anticancer, and anti-allergic activities (Di Carlo et al. 1999; Montoro et al. 2005). Flavonoids, have been shown to be highly effective scavengers of most oxidizing molecules, including singlet oxygen and various free radicals implicated in several diseases (Nabayi et al. 2008).

As *S. mombin* fruit juice exhibited significant levels of phytochemicals, which make it a potent source of antioxidant agents. This fruits can be vital help especially in such rural and urban areas where modern health facilities are not accessible due to poor state of living. Recent studies have also shown the advantages of natural antioxidants in foods, as opposed to synthetic additives (Shalini 2012).

Physicochemical properties of the fruit juice of *S. mombin*

The most important physicochemical properties of the crude *S. mombin* fruit juice were also determined. These values are displayed in Table 2.

The nutritional and energy values of foods are known through proximate analysis, but this gives a certain insight to the food quality. Earlier, Adepuju (2009) reported higher values for Nigerian *S. mombin* fruit juice (g/100 g) in moisture (82.3), lipid (2.0), crude fibre (4.2) and ash (1.0) than our values. From Brazil, Tiburski and co-workers (2011) reported moisture (83.66), fat (0.62), fiber (1.87), ash (0.76), total acidity (20.85), protein (1.06), carbohydrate (13.90) and pH (2.83) values, which are not significantly different from those we measured in this study.

Mineral composition of the crude fruit juice of *S. mombin*

Ten trace elements (potassium, manganese, copper, sodium, zinc, magnesium, lead, phosphorus, calcium, and iron) were determined from the crude fruit juice of *S. mombin*. The measured concentrations are displayed in Table 3. Some of these values are higher than what are reported for apricots, apple, banana, blackberry, grape fruit, cherries, lemon, lime, pear, pineapple, watermelon and other common and frequently consumed fruits (Cunningham et al. 2001; Emsley 2001; Decuyper 2005).

Minerals or trace elements are either essential or non-essential, depending on whether or not they are required for human nutrition and have metabolic roles in the body (Reilly 2002). Fruits generally have in their composition a great variety of vitamins and essential minerals, which makes them beneficial to our diet. In this study, low levels of manganese, iron, sodium, copper and lead were observed in the crude *S. mombin* fruit juice though higher content of magnesium, phosphorus, potassium and zinc have been reported present in some other edible fruits. Albino et al. (1999), described *S. mombin* as a fruit with high content of potassium, along with jackfruit, soursop, jenipapo (*Genipa americana*) and mangaba (*Hancornia speciosa*). The phosphorus content of *S. mombin* as found in this study is one of the highest among the commonly consumed fruits with levels close to those of Ceriguela (*Spondias purpurea*), Pequi (*Caryocar brasiliense*) and Passion fruit (NEPA-UNICAMP 2006).

Same mineral contents evaluated from *S. mombin* fruit juice by Adepoju (2009), cannot be compared because of the different methods used in analysis. Though several researchers have evaluated and reported different mineral contents of *S. mombin*, the differences can be attributed to the origin of the fruit. It may be dependent on the environmental conditions of each region (Leterme et al. 2006), since the minerals are absorbed from the soil. Furthermore, genetic factors, the use of fertilizers (Sanchez-Castillo et al. 1998) and the maturity stage of the tree that produces the fruits could also affect these results. The mineral contents of sodium, magnesium, potassium and phosphorus reported by Tiburski et al. (2011) from Brazil, were a bit higher than what was obtained in our

Table 2. Physicochemical properties of the fruit juice of *S. mombin*.

Physicochemical properties	Mean \pm SD
pH	3.63 \pm 0.01
Moisture (g/100 g)	80.66 \pm 0.02
Protein (g/100 g)	6.03 \pm 0.04
Fiber (g/100 g)	0.64 \pm 0.02
Ash (g/100 g)	0.54 \pm 0.03
Fat (g/100 g)	1.85 \pm 0.02
Total titratable acidity	28.52 \pm 0.04
Carbohydrate (g/100 g)	11.61 \pm 0.04

Table 3. Mineral content of the crude fruit juice of *S. mombin*.

Minerals	Mean \pm SD (mg/100 g)
Sodium (Na)	4.45 \pm 1.42
Magnesium (Mg)	12.36 \pm 0.03
Phosphorous (P)	28.22 \pm 1.02
Potassium (K)	276.27 \pm 2.65
Calcium (Ca)	13.04 \pm 0.53
Manganese (Mn)	0.15 \pm 0.01
Iron (Fe)	0.48 \pm 0.01
Copper (Cu)	0.121 \pm 0.32
Lead (Pb)	0.01 \pm 0.00
Zinc (Zn)	136.42 \pm 1.35

study, while we measured higher calcium, manganese, iron, copper, lead, and zinc levels.

Conclusions

The level of antioxidant in fresh fruits are in wholesome than the preserved in the commercial fruit juices. Many of these compounds are heat sensitive, which are destroyed during the industrial processing of juices. Additionally, these antioxidants frequently used for preservative purposes and are synthetic additives in the product. We are of the opinion that consumption of fresh *S. mombin* fruit juice will promote a better health status as the antioxidants, physicochemicals and minerals analyzed are vital and valuable than some of the many recognized fruits. The analysis of the crude fruit juice of *S. mombin* in this study is in support of the several reports of health benefits and medicinal value of this edible fruit.

Acknowledgement

We acknowledged all the Laboratory Technologists of the Biological Sciences Department of Afe Babalola Univer-

sity (Ado Ekiti, Nigeria) for their assistance in some of the analysis.

References

- Aberoumand A, Deokule SS (2010) Elements evaluation of some edible vegetables and fruits of Iran and India. *Asian J Agric Sci* 2(1):35-37.
- Adepoju OT (2009) Proximate composition and micronutrient potentials of three locally available wild fruits in Nigeria. *Afr J Agric Res* 4(9):887-892.
- Adewale A, Olubukola SO, Olusoji AO (2015) Nutritive values and antioxidant activity of *Citrullus lanatus* fruit extract. *Food Nutri Sci* 6:1056-1064.
- Akhtar S, Naz S, Tuseef SM, Mahmood S, Nasir M, Ahmad A (2010) Physicochemical attributes and heavy metal content of mangoes (*Mangifera indica*) cultivated in different regions of Pakistan. *Pak J Bot* 42(4):2691-2702.
- Albino E, Barreto RLSB, Coelho ACD, Coelho RD, Mendes ACR (1999) Análise da concentração de potássio em alguns frutos cultivados no nordeste. *Higiene Alimentar* 13(62):34-36. [in Portuguese]
- Ali L, Alsanius BW, Rosberg AK, Svensson B, Nielsen T, Olsson ME (2011) Effects of nutrition strategy on the levels of nutrients and bioactive compounds in blackberries. *Eur Food Res Technol* 234:33-34.
- Ali SS, Kasoju N, Luthra A, Singh A, Sharanabasava H, Sahu A (2008) Indian medicinal herbs as sources of antioxidants. *Food Res Int* 41(1):1-15.
- AOAC Methods of analysis of AOAC (2005) International (18th ed.). Arlington VA: Association of Official Analytical Chemists.
- Awogbindin IO, Tade OG, Metibemu SD, Olorunsogo OO, Farombi EO (2014) Assessment of flavonoid content free radical scavenging and hepatoprotective activities of *Ocimum gratissimum* and *Spondias mombin* in rats treated with dimethylnitrosamine. *Arch Bas App Med* 2:45-54.
- Benzie IF, Strain JJ (1996) The ferric reducing ability of plasma (FRAP) as a measure of "Antioxidant Power": The FRAP assay. *Anal Biochem* 39:70-76.
- Cavin A, Hostettmann K, Dyatmyko W, Potterat O (1998) Antioxidant and lipophilic constituents of *Tinospora crispa*. *Planta Med* 64:393-396.
- Cunningham J.H., Milligan G., Trevisan L (2001) Minerals in Australian fruits and vegetables - a comparison of levels between the 1980s and 2000. Food Standards Australia New Zealand, Canberra.
- Decuyper Dr. (2005) Nutrient contents of fruits, nuts and vegetables. *Health Alternative* 2000.com: 1-6.
- Di Carlo G, Mascolo N, Izzo AA, Capasso F (1999) Flavonoids: old and new aspects of a class of natural therapeutic drugs. *Life Sci* 65:337-353.
- Emsley J (2001) *Nature's Building Blocks: An A-Z Guide to the Elements*. Oxford University Press, New York
- Famobuwa EO, Oloyede H, Amuho U, Adekunbi A (2016a) In-vitro antioxidant and antibacterial properties of an anti-inflammatory herbal mixture of *C. jagus* L. (Amaryllidaceae) *E. hirta* Linn (Euphorbiaceae) and *T. tetraptera* Taub (Mimosaceae). *Eur J Pharm Med Res* 3(2):50-53.
- Famobuwa EO, Osho IB, Akinlami OO, Agbowuro AA (2016b) Anti-inflammatory activities of the chloroform extract of the fruit of *Tetracarpidium conophorum* (Mull. Arg.) (Nigerian Walnuts). *J Adv Med Pharm Sci* 6(1):1-7.
- Jensen GS, Wu X, Patterson KM, Barnes J, Carter SG, Scherwitz L, Beaman R, Endres JR, Schauss AG (2008) In vitro and in vivo antioxidant and anti-inflammatory capacities of an antioxidant-rich fruit and berry juice blend. Results of a pilot and randomized, double-blinded, placebo-controlled, crossover study. *J Agric Food Chem* 56(18):8326-8333.
- Kaur C, Kapoor HC (2001) Antioxidants in fruits and vegetables - the millennium's health. *Int J Food Sci Technol* 36:703-725.
- Leterme P, Buldgen A, Estrada F, Londoño AM (2006) Mineral content of tropical fruits and unconventional foods of the Andes and the rain forest of Colombia. *Food Chem* 95(4):644-652.
- Lu M, Yuan B, Zeng M, Chen J (2011) Antioxidant capacity and major phenolic compounds of spices commonly consumed in China. *Food Res Int* 44(2):530-536.
- Montoro P, Braca A, Pizza C, De Tommasi N (2005) Structure-antioxidant activity relationships of flavonoids isolated from different plant species. *Food Chem* 92:349-355.
- Nabavi SM, Ebrahimzadeh MA, Nabavi SF, Hamidinia A, Bekhradnia AR (2008) Determination of antioxidant activity, phenol and flavonoid content of *Parrotia persica* mey. *Pharmacol online* 2:560-567.
- NEPA-UNICAMP (2006) Tabela brasileira de composição de alimentos (2nd ed.) Campinas: Núcleo de Estudos e Pesquisas em Alimentação-UNICAMP. [in Portuguese]
- Reilly R (2002) Minerals. In CJK Henry and C Chapman (Eds) *The Nutrition Handbook for Food Processors*. CRC Press, Boca Raton, FL, USA. 97-116.
- Sakanaka S, Tachibana Y, Okada Y (2005) Preparation and antioxidant properties of extracts of Japanese persimmon leaf tea (kakinoha-cha). *Food Chem* 89:569-575.
- Sahreen S, Khan MR, Khan RA (2014) Comprehensive assessment of phenolic and antiradical potential of *Rumex hastatus* D. Don. roots. *BMC Complement Altern Med* 14:47.
- Sanchez-Castillo CP, Dewey PJS, Aguirre A, Lara JJ, Vaca R, Leon de La Barra P, Ortiz M, Escamilla I, James WPT (1998) The mineral content of Mexican fruits and veg-

- etables. *J Food Comp Anal* 11(4):340-356.
- Schinella G, Fantinelli JC, Tournier H, Prieto JM, Spegazzini E, Debenedetti S, Mosca SM (2009) Antioxidant and cardioprotective effects of *Ilex brasiliensis*: A comparative study with *Ilex paraguariensis* (yerba mate). *Food Res Int* 42:1403-1409.
- Shalini S (2012) Natural antioxidants - A Review. *Int J Phytother* 2(1):7-15.
- Singleton VL, Rossi JA (1965) Colorimetry of total phenolics with phospho-molybdic-phosphotungstic acid reagents. *Am J Enol Viticul* 16:144-158.
- Sreeramulu D, Raghunath M (2010). Antioxidant activity and phenolic content of roots, tubers and vegetables commonly consumed in India. *Food Res Int* 43(4):1017-1020.
- Tadhani MB, Patel VH, Subhash R (2007) *In vitro* antioxidant activities of *Stevia rebaudiana* leaves and callus. *J Food Comp Anal* 20(3-4):323-329.
- Temple NJ (2000) Antioxidants and disease: more questions than answers. *Nutr Res* 20:449-59.
- Tiburski JH, Amauri R, Rosires D, Ronoel L de Oliveira G, Sidney P (2011) Nutritional properties of yellow mombin (*Spondias mombin* L.) pulp. *Food Res Int* 44: 2326-2331.
- Vasco C, Ruales J, Kamal-Eldin A (2008) Total phenolic compounds and antioxidant capacities of major fruits from Ecuador. *Food Chem* 111:4816-4823.
- Willett WC (1994) Diet and health: what should we eat? *Science* 254:532-537.
- Xin SZ, Song YJ, Lv C, Rui YK, Zhang FS, Xu W, Wu D, Wu S, Zhong J, Chen DL, Chen Q, Peng FT (2009) Application of synchrotron radiation X-ray fluorescence to investigate the distribution of mineral elements in different organs of greenhouse spinach. *Hort Sci* 36 (4):133-139.
- Yazdanparast R, Ardestani A (2007) In vitro antioxidant and free radical scavenging activity of *Cyperus rotundus*. *J Med Food* 10:667-674.