

ARTICLE

Physiological and morphological responses of sugar beet (*Beta vulgaris* L.) subjected to nano-boron oxide at different growth stages

Alireza Pirzad*, Mahmood Mazlomi Mamyandi, Razieh Khalilzadeh

¹ Department of Plant Production and Genetics, Faculty of Agriculture and Natural Resources, Urmia University, Urmia, Iran

ABSTRACT The deficiency and toxicity of boron may lead to noticeable yield reduction and quality loss of sugar beet. To evaluate the effects of nano-boron on the root yield and quality of sugar beet, factorial experiment was conducted based on a randomized complete block design with three replications at Urmia University. Treatments were spraying of nano-boron oxide (0, 2, 3 and 4 g L⁻¹) at different growth stages (20, 40, 60, 80 and 100% of ground cover). Increasing levels of boron up to 3 g L⁻¹ resulted in the highest SPAD (Soil Plant Analysis Development) value, leaf number and relative water content. With optimal leaf area (86.47 cm²), the highest yields of root, sugar, white sugar, and technological sugar (144.53, 28.23, 26.19 and 25.32 t ha⁻¹) were determined in the treatment of 4 g L⁻¹ boron at 40% of ground covered. Increasing nano-boron level under different growth stages increased sugar and white sugar contents, on contrary impurities (Na, K and α-amino-N) loss and molasses sugar percentage were decreased. More than 3 g L⁻¹ nano-boron exhibited the highest values of purity. Application of 3 g L⁻¹ boron at 40% of ground cover increased sugar and white sugar contents by 12.45 and 18.72%, respectively.

Acta Biol Szeged 63(2):103-111 (2019)

KEY WORDS

growth stage
impurity
sugar beet
molasses sugar
nano-boron

ARTICLE INFORMATION

Submitted

18 December 2019.

Accepted

03 February 2020.

*Corresponding author

E-mail: a.pirzad@urmia.ac.ir

Introduction

Sugar beet (*Beta vulgaris* L., Amaranthaceae) is one of the important sugar crops with high yield, high adaptability and high biological activity. It is also rich in minerals and organic nutrients (Grzegorzewski et al. 2017). Although sugar beet is well adapted to a wide range of growing conditions and soils, nutritional disorders caused by boron deficiency are quite common (Dordas 2007). Sustainable production requires the efficient use of inputs including adequate and balanced fertilization of both macro- and micro-nutrients (Singh et al. 2017). Deficiency of soil nutrients such as nitrogen (N), phosphorus (P), potassium (K), zinc (Zn) and boron (B) should be added to the rhizosphere according to plant needs and has been known as the major limitations in beet crop production (Abidow 2012). Boron is unique among the essential mineral micronutrients because of normal presence in soil solution as a non-ionized molecule over the vast pH range (Hirparan et al. 2018). It is also essential for plant in development and growth (Abido 2012) through the cell wall structure, membrane integrity and function (Nyomora et al. 2019), sugar translocation from source to sink

(Rawashdeh and Sala 2013) physiological functions such as carbohydrates metabolism, indole acetic acid metabolism, formation amino acid (Singh et al. 2017), nitrogen fixation (Rawashdeh and Sala 2013) and photosynthetic pigments (Abd El-hady 2017).

Deficiency of boron causes reduction in photosynthesis due to disturbs the activities of proton pumping ATPase and electron transport chain (Nadeem et al. 2019; Rehman et al. 2018), inhibition of leaf expansion, inhibitions root elongation through limiting mitosis and cell enlargement and division (Gemici et al. 2003), early enlargement due to clear limitation of its phloem mobility and reduced growth of new shoots and leaf (Ullah et al. 2013; Ali et al. 2015). Without an adequate supply and consumption of boron in large quantities, it may lead to marked yield reduction and quality loss of sugar production in soil application (Abbas et al. 2014; Tlili et al. 2017; Armin and Asgharipour 2012). Because boron has been considered to have only limited phloem mobility and cannot readily be redistributed within the plant (Brown et al. 1999), the amount of yield losses directly depends upon the duration of deficiency and the plant growth stage at which it occurs (Ali et al. 2015). Deficiencies of boron result in many anatomical, biochemical and physiological changes

in plants (Ali et al. 2015). Therefore, the management of boron in soil has become a worldwide agricultural problem in the recent years (Tlili et al. 2017).

Foliar fertilization has the advantage of low application rates, treated rapidly, uniform distribution and quick plant responses to applied nutrients (Asad et al. 2003; Saadati et al. 2013). Moreover, a number of previous studies have increased the significance of the role of foliar boron application in the productivity of crop plants (Perica et al. 2001; Dordas 2006; Abido 2012; Kristek et al. 2006). Also, to get the desired results, nanomaterial can be utilized by foliar application with much-decreased concentration (Prasad et al. 2012). After entering the cells the nanoparticles transport from one cell to another through plasmodesmata. The chemical and biological activities of most substances increase at the nanoscale (Dewdar et al. 2018). Root yield, sugar percentage significantly increased by increasing boron doses (Abbas et al. 2014). Dordas et al. (2006) reported that spraying of boron lead to a higher quality of sugar and root yield compared to the time using boric acid mixed with soil. El-Geddawy and Makhlof (2015) found that there was a significant positive increase in root diameter and root length of sugar beet due to the gradual increase in the spraying concentration of boron from 105 to 210 ppm. Armin and Asgharipour (2012) reported that the maximum root yield and sugar percentage was achieved by foliar application of 12% boric acid. Abd El-hady (2017) reported that root and sugar yields were increased by 19.4% and 39.5% compared with control treatment.

A better understanding of the physiological basis of the response of sugar beet may help in programs aiming to evaluate yield. Therefore, this work aimed to evaluate the effects of different amounts and time of nano-boron oxide spraying on the quantitative and qualitative aspects of sugar beet.

Materials and Methods

Site and experimentation

All experiments were conducted during 2011-2012 cropping seasons in North-Western Iran Iran, i.e. Naghadeh (27°45' N latitude and 22°37' E longitude; Alt 1300 m) and were situated in the wet zone with moderate winter and hot summer. The experimental design was randomized factorial experiment based on a randomized complete block design with three replications. The soil type was silty loam and possessed around 7.95 pH, EC about 2.3 dS m⁻¹, total organic C = 1.20% and Zn = 32 mg kg⁻¹. The experimental soil was fertilized with 250 kg N ha⁻¹ in the form of urea (was applied as ½ at sowing, ½ at 6-8 leaf), 250 kg P ha⁻¹ in the form of triple superphosphate, and

100 kg K ha⁻¹ in the potassium nitrate at planting time. Experiment factors were the amount of nano-boron (Nano-B) oxide concentration (0, 2, 3 and 4 g L⁻¹ of nano chelate powder, with 99.5% of purity and 80 nm particle size, obtained from Khazra Company containing 9% chelated boron, absorbable at pH 3-11, and completely soluble in water) and spraying time included (20, 40, 60, 80 and 100 of ground cover by plant canopy). Each plot was consisted of 5 rows with 5 m long. The inter row and intra-row spacing was 10 and 15 cm, respectively. The sugar beet cultivar (Montarosa cv. a commonly grown cultivar of sugar beet in Naghdeh area) was sown at the depth of 20 cm on May 10.

Measurements of quantity and quality parameters of sugar beet

Harvesting was done manually 180 days after sowing (DAS). In order to measure the root length (cm), root diameter (cm) and leaf number traits, 10 plants in each plot were randomly harvested. Root yield (t ha⁻¹) was obtained from plants harvested of 5 m² in each experimental unit and juice quality characteristics were analyzed. The percentage of sucrose was determined according to Le-Docte (1927). Sodium and potassium (%) were determined by using a flame photometer (Model 410 Classic), nitrogen was determined according to the semi-micro Kjeldahl method (Model NA 1500) (Edwards 2014). Total soluble solids (TSS%) was measured in fresh roots using hand refractometer (model REF-113ATC). Juice purity% was also determined as a ratio between sucrose% and TSS% according to Carruthers and Oldfield (1961).

Sugar yield and root quality were calculated via the following equations:

$$SY (t ha^{-1}) = RY (t ha^{-1}) \times SC(\%)$$

Where, SY: sugar yield; RY: root yield; SC: sugar content.

$$MS = 0.12 (K + Na) + 0.24(\alpha\text{-amino-N}) + 0.48$$

Where, MS: molasses sugar (%); K: potassium (mmol/100 g root), Na: sodium (mmol/100 g root); α -amino-N: alpha-amino-nitrogen (mmol/100 g root) (Buchholz et al. 1995).

$$AC: (K + Na)/ N.$$

Where, AC: alkalinity coefficient.

$$WSC(\%) = SC\% - MS\%$$

Where, WSC: white sugar content.

$$WSY (t ha^{-1}) = RY \times WSC \%$$

Where, WSY: white sugar yield.

$$\text{TSY (t ha}^{-1}\text{)} = \text{RY (t ha}^{-1}\text{)} \times [\text{SC (\%)} - \text{loss of sugar productivity (\%)}]$$

Where, TSY: technological sugar yield (Buchholz et al. 1995).

$$\text{LSP (\%)} = \text{MS (\%)} + 0.6$$

Where, LSP: loss of sugar productivity (Carruthers et al. 1961).

$$\text{Sugar productivity (SP)} = \frac{\text{SC} - \text{loss of sugar productivity}}{\text{SC}}$$

Where, SP: sugar productivity.

The leaf length was measured as the distance between the beginning of leaf formation on the leaf stem and the top of the leaf. The leaf width was measured at its widest point with a ruler. Based on measured leaf width (W, mm) and leaf length (L, mm) the area of each leaf (A, mm²) is calculated using the following relationship (Mirschel 2018):

$$A = W \times L \times 0.675$$

The chlorophyll content of leaves was estimated with a SPAD-502 (Konica Minolta Sensing, Osaka, Japan) (Jifon et al. 2019). Relative water content was estimated according to the method of Tambussi et al. (2005).

Analysis of variance (ANOVA) and means comparison on data was performed using the SAS Statistical Package Program. Least Significant Difference (LSD) method was used to test the differences between means comparison of main effects and interactions.

Results

Analysis of variance showed that the significant interaction effect between Nano-B concentration and spraying stage on the relative water content (RWC), leaf area, root length, root and sugar yield, white and technological sugar yield, sugar content, white sugar content, sugar productivity, loss of sugar productivity, sodium, potassium, α -amino-N and molasses sugar. There was a significant effect of Nano-B foliar application on the alkalinity coefficient (Table 1). SPAD, leaf number, root diameter, purity% and TSS% were affected by the Nano-B concentration and spraying stage (Table 1).

Chlorophyll index, relative water content, leaf area and leaf number

The maximum of SPAD (73.50) and leaf number (52.26) was obtained in B₃ and B₄, respectively (Table 1). Chlorophyll index was increased about by 34.71% in 3 g L⁻¹ Nano-B (Table 1). The highest content of SPAD and leaf number markedly increased from spraying of Nano-B at 80 of ground cover as G₄ (Table 1). However, the difference in mention traits was not statistically significant in a comparison between G₂, G₃ and G₄. The increasing of Nano-B at all growing stages up to 3 g L⁻¹ resulted in the highest RWC. The highest and lowest RWC was respectively achieved in B₃G₂ and B₁G₅ (Table 2). A gradual increase in leaf area as growth stages improved up to G₄ was recorded regardless of boron levels. The application of B₃G₄ caused an increase in leaf area of 60% in comparison with B₁G₄.

Root length and diameter

It could be noticed that increasing boron rates significantly increased root diameter. The plant sprayed with 3 g L⁻¹ of nano-boric acid revealed the highest root diameter (12.70 cm). Data also cleared that the late application of boron (G₄) recorded the highest value of root diameter (12.79 cm). Results showed that, the crops were fertilized early or late at different rates of boron, had any considerable differences on root length. Application of 2 g L⁻¹ at 100% of ground covered (B₂G₅), produced the highest root length (38.16 cm), while the lowest value (25.50 cm) was recorded in the B₁G₁ (Table 2).

Sugar beet yields

Application of 4 g boron L⁻¹ at the early stage of plant (40% ground cover), can significantly increase root yield (144.53 t ha⁻¹), sugar yield (28.23 t ha⁻¹), white sugar yield (26.19 t ha⁻¹), and technological sugar yield (25.32 t ha⁻¹) to the highest amounts. Spraying nano-boric acid at the levels of B₂, B₃ and B₄ increased root yield by about 29%, 48% and 79% at early growth stage (G₂), as compared to control treatment (B₁), respectively. However, Abd El-hady (2017) reported that B element (1.0 kg B/ha) was increased root and sugar yields by 19.4% and 39.5% compared with control treatment.

Quality of sugar beet

The highest sugar and white sugar content were found to be in the B₃G₂ treatment with average of 19.86% and 18.39%, respectively. However, their effects were also similar to B₄G₂. Also, the lowest of SC and WSC are related to B₁G₁ treatment with an average of 15.80% and 14.14%, respectively (Table 3). Compared to the control, spraying 3 g L⁻¹ Nano-B at 40% of ground covered improved SC and WSC by 12.45% and 15.37%, respectively. Data in Table 1

Table 1. Effects of nano-boron oxide concentration and spraying time (growth stage) on growth and yield of sugar beet.

Nano-B (B)	SPAD	RWC (%)	Leaf number	Leaf area (mm ²)	Root length (cm)	Root diameter (cm)	Root yield	Sugar yield		TSY
								(t ha ⁻¹)		
B ₁ = no nano-B as control	54.56b	73.98b	33.73c	49.52c	26.70c	10.73b	88.77d	14.93d	13.39d	12.86d
B ₂ = 2 g L ⁻¹	59.55b	82.07a	41.46b	67.63b	30.53b	11.23b	108.25c	18.92c	17.26c	16.61c
B ₃ = 3 g L ⁻¹	73.50a	84.82a	52.26a	74.55ab	31.00b	11.36b	117.09b	21.73b	20.05b	19.35b
B ₄ = 4 g L ⁻¹	68.35a	83.27a	49.33a	82.85a	34.68a	12.70a	127.65a	24.15a	22.42a	21.65a
LSD (p<0.05)	5.20	4.08	4.96	10.47	2.49	1.24	8.05	1.70	1.60	1.55
Growth stage (G)										
G ₁ = 20% of ground cover	58.89b	81.74ab	41.58b	55.65c	28.87cd	10.25c	98.94b	17.28b	15.82b	15.23b
G ₂ = 40% of ground cover	64.38a	83.48a	44.16ab	69.57abc	27.91d	11.04bc	114.39a	21.36a	19.65a	18.96a
G ₃ = 60% of ground cover	65.70a	81.59ab	46.16ab	73.80ab	30.26bc	11.26bc	113.007ab	20.69a	19.09a	18.41a
G ₄ = 80% of ground cover	66.92a	81.68ab	48.08a	80.15a	31.20b	12.79a	115.52a	20.40ab	18.66ab	17.96ab
G ₅ = 100% of ground cover	64.07ab	76.69b	41b	64.03bc	35.37a	12.18ab	110.35ab	19.94ab	18.19ab	17.53ab
LSD (p<0.05)	5.36	5.53	5.30	14.18	2.10	1.33	14.49	3.33	3.24	3.16
B × G	ns	**	ns	*	**	ns	**	*	**	*
C.V.	10.14	5.18	14.51	14.96	8.28	11.89	7.12	8.31	8.55	8.65

ns: show no significant differences.
 * and **: show significant differences at 0.05 and 0.01 probability level, respectively.
 RWC: relative water content; WSY: white sugar yield; TSY: technological sugar yield.

Table 1 (Continued). Effects of nano-boron oxide concentration and spraying time (growth stage) on growth and yield of sugar beet.

Nano-B (B)	SC (%)	WSC (%)	Purity %	SP	LSP	Na K α-amino-N			AC	MS	TSS %
						(mmol/100 g root)					
B ₁ = no nano-B as control	16.84b	15.11c	75.92c	0.861d	2.33a	1.20a	6.84a	1.18a	6.99c	1.73a	22.20a
B ₂ = 2 g L ⁻¹	17.50b	15.96b	82.83b	0.877c	2.13b	0.79b	6.27b	0.85b	8.46b	1.53b	21.15b
B ₃ = 3 g L ⁻¹	18.54a	17.11a	92.12a	0.890b	2.03c	0.68b	5.77c	0.74b	8.97b	1.43c	20.18bc
B ₄ = 4 g L ⁻¹	18.90a	17.54a	95.72a	0.896a	1.96d	0.52c	5.61c	0.60c	10.44a	1.36d	19.91c
LSD (p<0.05)	0.69	0.67	4.90	0.005	0.07	0.117	0.45	0.118	1.12	0.07	0.96
Growth stage (G)											
G ₁ = 20% of ground cover	17.40b	15.92b	88.42a	0.879b	2.08ab	0.69b	7.07abc	0.77a	9.00a	1.47ab	19.95c
G ₂ = 40% of ground cover	18.53a	17.009a	88.30a	0.884a	2.12ab	0.70b	6.30ab	0.84a	8.71a	1.52ab	21.09abc
G ₃ = 60% of ground cover	18.21ab	16.77ab	85.26ab	0.886a	2.044b	0.82a	5.61c	0.80a	8.73a	1.44b	21.43ab
G ₄ = 80% of ground cover	17.58ab	16.06ab	81.89b	0.878b	2.11ab	0.90a	6.00bc	0.87a	8.44a	1.51ab	21.54a
G ₅ = 100% of ground cover	18.01	16.41ab	89.38a	0.877b	2.20a	0.87a	6.63a	0.93a	8.70a	1.60a	20.30bc
LSD (p<0.05)	0.99	1.072	4.68	0.0043	0.137	0.094	0.59	0.22	1.65	0.137	1.22
B × G	**	*	ns	*	**	**	**	*	ns	**	ns
C.V.	4.06	4.42	6.54	0.58	2.63	14.23	6.56	16.43	17.11	3.66	5.20

ns: show no significant differences.
 * and **: show significant differences at 0.05 and 0.01 probability level, respectively.
 SC: sugar content; WSC: white sugar content; K: potassium; Na: sodium; α-amino-N: alpha-amino-nitrogen; AC: alkalinity coefficient; TSS: total soluble solids; SP: sugar productivity; LSP: loss of sugar productivity MS: molasses sugar.

noticeably showed that B₄ and B₃ treatments recorded the highest values of purity percentage by 95.72% and 92.12%, respectively. It is worth mentioning that, considerable differences in purity% were not significant (Table 1). The later application at G₅ insignificantly surpassed the earlier application at G₃ in effecting purity%. Increasing the doses of B application from 0 to 4 g L⁻¹ provided the lowest LSP

and MS 1.79 and 1.19 % with a decrease of 20.44% and 27.87%, respectively at the 60% of ground covered.

Root impurities (K, Na and α-amino-N) and AC

The values of impurities differed greatly due to the different treatments of time and boron rates (Table 3). There was a negative relationship between impurities and boron

Table 2. Means comparison the effects of Nano-B concentration and spraying time (growth stage) on RWC, leaf area, root length, root yield, sugar yield, WSY and TSY of sugar beet.

Treatment		RWC (%)	Leaf area (mm ²)	Root length (cm)	Root yield	Sugar yield (t ha ⁻¹)	WSY	TSY
Nano-B	Growth stage							
B ₁	G ₁	77.34±3.65	52.10±6.53	25.50±2.50	82.87±7.17	13.08±1.055	11.17±0.89	11.21±0.85
	G ₂	79.18±2.06	50.66±2.51	26.66±1.52	80.93±1.88	14.28±0.42	12.90±0.39	12.41±0.39
	G ₃	75.65±2.64	58.61±6.88	25.33±1.52	90.56±9.19	15.06±1.72	13.55±1.54	13.01±1.49
	G ₄	79.16±3.50	57.57±2.97	27.50±2.17	100.88±8.64	16.39±1.17	14.68±1.09	14.08±1.04
	G ₅	58.57±5.09	28.69±1.69	36.16±1.89	88.61±8.10	15.84±1.66	14.13±1.47	13.60±1.42
B ₂	G ₁	81.64±3.22	51.49±6.04	31.33±1.25	98.36±6.93	17.43±1.02	15.99±0.96	15.40±0.92
	G ₂	82.22±3.87	66.32±10.35	27.50±2.00	112.21±5.77	19.13±0.73	17.46±0.67	16.79±0.65
	G ₃	82.33±1.86	74.23±15	34.56±3.84	106.85±10.80	19.51±1.46	17.91±1.36	17.26±1.30
	G ₄	81.25±1.16	84.36±5.99	34.83±2.84	115.25±10.36	19.68±2.05	17.87±1.99	17.18±1.93
	G ₅	82.93±2.44	61.78±17.16	38.16±2.25	108.61±7.77	18.87±2.30	17.09±2.10	16.44±2.05
B ₃	G ₁	85.23±4.76	53.45±4.49	29.16±1.52	111.17±3.47	19.11±0.85	17.58±0.78	16.92±0.78
	G ₂	88.86±9.10	74.83±9.15	25.66±2.36	119.89±10.14	23.81±2.23	22.05±2.10	21.34±2.05
	G ₃	85.41±1.38	76.06±12.47	33.00±4.35	119.06±13.26	22.97±2.30	21.28±2.19	20.57±2.13
	G ₄	83.56±10.56	92.13±18.71	32.16±2.75	118.42±5.98	21.37±0.59	19.59±0.46	18.88±0.43
	G ₅	83.51±2.20	79.30±15.30	30.83±2.84	116.90±6.79	21.38±1.89	19.74±1.77	19.04±1.73
B ₄	G ₁	82.78±2.47	65.57±9.14	29.50±1.32	103.38±7.20	19.49±1.16	18.02±1.10	17.40±1.07
	G ₂	83.66±0.31	86.47±10.87	31.83±1.75	144.53±6.95	28.23±2.026	26.19±2.01	25.32±1.98
	G ₃	82.99±0.46	86.29±8.96	28.16±1.15	135.54±6.011	25.24±2.89	23.62±2.85	22.80±2.82
	G ₄	82.74±1.99	89.36±8.50	30.33±2.75	127.54±2.52	24.14±0.60	22.49±0.56	21.72±0.54
	G ₅	81.74±2.03	86.56±14.19	36.33±4.01	127.28±7.60	23.67±1.56	21.81±1.44	21.04±1.40
LSD _{0.05}		6.94	16.97	4.21	13.01	2.73	2.58	2.52

B₁ indicates no application; B₂, B₃ and B₄ indicate application of 2, 3 and 4 g L⁻¹ of Nano-B, respectively.

G₁, G₂, G₃, G₄ and G₅ indicate foliar application Nano-B at 20, 40, 60, 80 and 100% of ground cover, respectively.

foliar application. Data in Table 3 exposed that, increasing the boron concentrations from 0 to 4 g L⁻¹ contributed the last potassium and α -amino-N content of 4.58 and 0.45 mmol/100 g root at 60% growth stage. On the other hand, without the application of boron (B₁) resulted in the maximum mean values of K (7.69) and α -amino-N (1.47) content at B₁G₅. The significant lowest sodium content 0.50 and 0.49 mmol/100 g root related to a reduction in impurity 60.37% and 66.66%, at the application of B₄G₃ and B₄G₄, respectively.

Discussion

Foliar application of Nano-B at 3 g L⁻¹ resulted in a consistent improvement in vegetative growth of sugar beet, but on increasing Nano-B concentration up to B₄, vegetative growth decreased compared with B₃. In addition, the increases of yield-related responses like chlorophylls and leaf area (Table 2) of sugar beet at high Nano-B concentration (B₃ and B₄) could be reflected upon the increase of sugars percentage and reduction of impurities (Table 3), so, the optimal leaf area value for root and sugar yields

was 86.47 cm² at B₄G₂. Leaf number and low chlorophyll content at high boron concentration are associated with toxicity of this element (Armin and Asgharipour 2012). Adequate boron supply through foliar application improved the chlorophyll content, leaf number and leaf area enabling them to capture more light and produce more assimilate for loading to root. Ullah et al. (2013) reported that B deficiency causes to reduced growth of new leaf and shoot due to clear limitation of boric acid phloem mobility. Such enhancement effect of B could be related to the favorable influence of them on photosynthetic pigments (Wanas 2002; Abd El-hady et al. 2017), metabolism, enzyme activity (El-Sherbeny et al. 2007), photosynthesis efficiency (Abou El-Yazied and Mady 2012) which in turn encourage vegetative growth and increasing dry matter production. Also, this enhancement could be an indicator of expectable high sugar beet yield. Also, Abd El-hady et al. (2017) informed that these results might be attributed to that B is an essential element for photosynthetic pigments, where it increases CO₂ fixation, rates of photosynthetic O₂ evolution and decreases respiration and the activities of oxidative pentose phosphate enzymes.

It seems that the increase in root diameter at 3 g L⁻¹ of

Table 3. Means comparison the effects of Nano-B concentration and spraying time (growth stage) on sugar content, WSC, SP, loss of sugar productivity, Na, K, N and MS of sugar beet.

Treatment		SC (%)	WSC (%)	SP	LSP	Na	K	N	MS
Nano-B	Growth stage	(mmol/100g root)							
B ₁	G ₁	15.80±0.20	14.14±0.27	0.857±0.006	2.25±0.085	0.94±0.16	6.93±0.71	0.96±0.11	1.65±0.081
	G ₂	17.66±0.83	15.94±0.74	0.868±0.0025	2.32±0.094	0.96±0.27	7.03±0.87	1.17±0.24	1.72±0.096
	G ₃	16.63±0.73	14.97±0.75	0.863±0.0077	2.25±0.058	1.26±0.22	6.33±0.59	1.11±0.19	1.65±0.060
	G ₄	16.26±0.41	14.56±0.37	0.859±0.003	2.29±0.070	1.47±0.101	6.23±0.24	1.22±0.14	1.69±0.067
	G ₅	17.86±0.64	15.94±0.58	0.859±0.002	2.52±0.070	1.37±0.061	7.69±0.79	1.47±0.14	1.92±0.068
B ₂	G ₁	17.73±0.30	16.26±0.23	0.883±0.002	2.07±0.072	0.66±0.047	5.84±0.28	0.85±0.21	1.46±0.071
	G ₂	17.06±0.64	15.58±0.65	0.877±0.0049	2.08±0.011	0.76±0.035	5.93±0.13	0.84±0.03	1.48±0.011
	G ₃	18.30±0.79	16.79±0.77	0.885±0.0045	2.10±0.043	0.83±0.100	5.96±0.19	0.86±0.19	1.50±0.045
	G ₄	17.06±0.50	15.49±0.54	0.872±0.0073	2.17±0.081	0.81±0.081	9.56±0.32	0.86±0.14	1.57±0.081
	G ₅	17.33±0.90	15.69±0.83	0.871±0.0034	2.23±0.075	0.91±0.085	7.05±0.52	0.83±0.05	1.63±0.072
B ₃	G ₁	17.20±0.80	15.82±0.78	0.885±0.0049	1.97±0.055	0.63±0.062	5.59±0.30	0.60±0.09	1.37±0.057
	G ₂	19.86±0.80	18.39±0.79	0.896±0.0040	2.06±0.025	0.57±0.028	6.30±0.24	0.67±0.07	1.46±0.021
	G ₃	19.33±0.98	17.90±0.98	0.894±0.0061	2.02±0.050	0.69±0.060	5.56±0.27	0.80±0.20	1.42±0.052
	G ₄	18.06±0.50	16.56±0.50	0.883±0.0040	2.10±0.052	0.83±0.015	5.99±0.36	0.84±0.09	1.50±0.052
	G ₅	18.26±0.57	16.86±0.55	0.890±0.0023	2.00±0.20	0.67±0.105	5.41±0.21	0.78±0.12	1.39±0.019
B ₄	G ₁	18.86±0.64	17.44±0.66	0.893±0.0050	2.02±0.034	0.53±0.041	5.92±0.11	0.69±0.10	1.42±0.035
	G ₂	19.53±0.90	18.11±0.91	0.896±0.0050	2.02±0.052	0.52±0.058	5.94±0.48	0.68±0.03	1.41±0.054
	G ₃	18.60±1.56	17.40±1.58	0.903±0.0091	1.79±0.046	0.50±0.073	4.58±0.33	0.45±0.08	1.19±0.047
	G ₄	18.93±0.23	17.63±0.25	0.899±0.0030	1.90±0.043	0.49±0.072	5.23±0.20	0.55±0.18	1.30±0.043
	G ₅	18.60±0.52	17.13±0.55	0.888±0.0047	2.06±0.041	0.54±0.032	6.37±0.03	0.64±0.15	1.46±0.039
LSD _{0.05}		1.207	1.20	0.0085	0.092	0.188	0.66	0.22	0.091

B₁ indicates no application; B₂, B₃ and B₄ indicate application of 2, 3 and 4 g L⁻¹ of Nano-B, respectively. G₁, G₂, G₃, G₄ and G₅ indicate foliar application Nano-B at 20, 40, 60, 80 and 100% of ground cover, respectively.

Nano-B by means of high leaf number and chlorophyll content and efficient assimilates portioning towards sink parts. These results could be explained by the role of boron in plant metabolism, development and growth (Rawashdeh and Sala 2013; Abido 2012), cell wall formation and meristematic tissue extension and cell elongation the root (Nalini et al. 2013). Cell enlargement and increase in a number of cells contribute to the increase of yield can be due to the role of boron in the biosynthesis of auxin in the meristematic activity and increase in the IAA-oxidase activity. Similar observations were recorded by Abdelaal et al. (2015) and Dugger (1973) in sugar beet.

Armin and Asgharipour (2012) stated that the maximum root yield and sugar percentage was achieved by foliar application of 12% boric acid. Considering this, Nano-B spraying may be used to enhance root and sugar yield, resulting in reduce boron fertilizer. Foliar Nano-B application predominantly affects at vegetative growth compared with reproductive growth in sugar beet. Results in Table 2 exhibited that sugar yield and white sugar yield was significantly improved by increasing of B from B₁ to B₄. These results were true in the five growth stages. While that sugar content was decreased by B₄ when compared

with B₃ treated plants (Table 2), the increase in sugar yield accompanying high boron level might have been due to the increase in root yield as well as sucrose content. These results are in agreement by those of Gezgin et al. (2001).

It seems that, better translocation of photosynthates from high leaf area (Table 2) and higher dry matter accumulation with high root length led to increasing in root yield (Table 2). The increase in tops and roots fresh and dry weights, caused by boron application, could be attributed to the stimulating effect of boron on the photosynthesis process in the plants such as translocation of sugar and carbohydrates of assimilates from the top to root, which leads to increase in root and sugar yield. On the other hand, when photosynthetic activity is high, any factor that increases the leaf area may have a positive effect on WSY. The enhancement of dry matter in sugar beet roots may be attributed to the improvement of leaf area, leaf number, RWC and chlorophyll content which results in improvement of growth-related traits such as root length and root diameter, and consequently root yield sugar yield (Table 2). Similar results were also observed by Abdel-Motagally (2015) and Mohammadian et al. (2014) who reported that early beginning of photosynthetic

transmission from leaf to root and consequently, it would increase WSY at the harvest time.

The rise of WSC may be due to the increase in sugar percentage and the reduction of impurities in terms of sodium, potassium and α -amino-N content (Table 3). Further, if optimum B is available at the early growth stage, the plant continues to partition it in the sink part as have been observed in this study. This finding suggested that sugar content is the key factor conferring B phloem mobility due to the bonding of sugar and boric acid (Liakopoulos et al. 2005). The Nano-boron facilitates the transport of sugars in the plants because it had a crucial role in the biosynthesis of auxin (Dugger 1973; Ullah et al. 2013). The high amount of juice purity would be desirable to provide sugar content. The important role of the boron element on the percentage of purity comes through its beneficial effect on the values of sucrose content (Table 3). Boron dominates in the early-stage, building up the highest TSS for the sugar beets (Table 1).

The result showed that, at low concentration of boron, a lower percentage of sugar had been achieved. Therefore, sugar in the form of molasses in these plants is higher in amount than that which has received enough boron (Table 3). Reduction of impurities like sodium, potassium and α -amino-N content in beet roots cause to decrease of sucrose molasses under application of a higher rate of boron application. These results are an agreement with the finding of Abbas et al. (2014) who reported that increasing the concentration of B cause to the reduction of sugar molasses. Hellal et al. (2009) showed that juice purity, sugar and root yield of sugar beet improved by increasing of B spraying which could be related to the reduction of sodium and potassium uptake in root juice.

The reasons for the improvement in sugar beet quality could be due to the fact that B plays a role in cell division, enhanced enzymatic activity, the membrane integrity, calcium uptake and carbohydrate metabolism (Nalini et al. 2013; Rawashdeh and Sala 2013). An increased boron supply decreases the nitrate levels via inhibiting transcript level in the roots and altering the nitrate transporter activity, leading to reduced plasma membrane enzymes activities (Camacho-Cristobal and Gonzalez-Fontes 2007). Evidence proposes that sugar alcohols synthesis and the later transport of the B-sugar alcohol compound in the phloem to sink tissues is the main factor that confers phloem B mobility to a plant species (Brown et al. 1999). The least accumulation of sodium at the later application (G_3 and G_4) could be due to the increased of leaf area at this time and facilitate the improvement of B absorption. Similar results were obtained by Abbas et al. (2014) who reported that spraying dates lead to significantly different in sodium content. Also, Armin and Asgharipour (2012) and Abbas et al. (2014) showed that boron application

improved juice quality by declining K and Na content.

Conclusion

Application of Nano-B rates showed a significant increase in quantitative and qualitative sugar beet traits under study. The highest SPAD and RWC, leaf area and leaf number were observed in 3 g L⁻¹ of Nano-B resulted in a consistent improvement in vegetative growth of sugar beet but with increasing concentration of Nano-B mentioned parameters were decreased. The increasing of boron fertilizer at all growing stages resulting in the highest root yield, sugar content and white sugar content thus led to increasing sugar yield, white and technological sugar yield. The decrease in sucrose molasses in a high level of boron accompanying due to the reduction in impurities in terms of sodium, potassium and α -amino-N content in sugar beet roots. Therefore, B₄G₂ treatment (4 g L⁻¹ boron at 40% of ground covered) with the highest root and sugar yield may be recommended for the cultivation of sugar beet in terms of time and fertilizer saving.

Acknowledgements

We thank "Iran's National Elites Foundation" for their help during the course of experimentation.

References

- Abbas MS, Dewdar MDH, Gaber EI, Abd El-Aleem HA (2014) Impact of boron foliar application on quantity and quality traits of sugar beet (*Beta vulgaris* L.) in Egypt. Res J Pharm Biol Chem Sci 5(5):142-151.
- Abdelaal KAA, Shima AB, Shahrzd MMN (2015) Effect of foliar application of microelements and potassium levels on growth, physiological and quality characters of sugar beet (*Beta vulgaris* L.) under newly reclaimed soils. Mansoura J Plant Prod 6:123-133.
- Abdel-Motagally FMF (2015) Effect concentration and spraying time of boron on yield and quality traits of sugar beet grown in newly reclaimed soil conditions. ASSIUT J Agric Sci 46(6):15-26.
- Abido WAE (2012) Sugar beet productivity as affected by foliar spraying with methanol and boron. Int J Agr Sci 4(7):287-292.
- Abou El-Yazied A, Mady MA (2012) Effect of boron and yeast extract foliar application on growth, pod setting and both green pod and seed yield of broad bean (*Vicia faba* L.). J Appl Sci Res 8(2):1240-1251.
- Ali F, Ali A, Gul H, Sharif M, Sadiq A, Ahmed A, Ullah A,

- Mahar A, Kalhoro SA (2015) Effect of boron soil application on nutrients efficiency in tobacco lLeaf. *Am J Plant Sci* 6:1391-1400.
- Amin GA, Badr EA, Afifi MHM (2013) Root yield and quality of sugar beet (*Beta vulgaris* L.) in response to bio fertilizer and foliar application with micro-nutrients. *World Appl Sci J* 27(11):1385-1389.
- Armin M, Asgharipour M (2012) Effect of time and concentration of boron foliar application on yield and quality of sugar beet. *Am Eurasian J Agric Environ Sci* 12(4):444-448.
- Asad A, Blamey FPC, Edwards DG (2003) Effects of boron foliar applications on vegetative and reproductive growth of sunflower. *Ann Bot* 92(4):565-570.
- Brown PH, Bellaloui N, Hu H, Dandekar A (1999) Transgenically enhanced sorbitol synthesis facilitates phloem boron transport and increases tolerance of tobacco to boron deficiency. *Plant Physiol* 119(1):17-20.
- Buchholz K, Märländer B, Puke H, Glattkowski H, Thielecke K (1995) Neubewertung des technischen Wertes von Zuckerrüben. *Zuckerindustry* 120:113-121.
- Carruthers N, Oldfield JFT (1961) Methods for assessment of beet quality. *Int Sugar J* 63:137-139.
- Dewdar DHM, Abbas MS, El Hassanin AS, Abd El-Aleem HA (2018) Effect of nano micronutrients and nitrogen foliar applications on sugar beet (*Beta vulgaris* L.) of quantity and quality traits in marginal soils in Egypt. *Int J Curr Microbiol Appl Sci* 7(8):4490-4498.
- Dewdar MDH, Abbas MS, Gaber EI, Abd El-Aleem HA (2015) Influence of time addition and rates of boron foliar application on growth, quality and yield traits of sugar beet. *Int J Curr Microbiol Appl Sci* 4(2):231-238.
- Dordas C, Apostolides GE, Goundra O (2007) Boron application affects seed yield and seed quality of sugar beets. *J Agric Sci* 145:377-384.
- Dordas C (2006) Foliar boron application affects lint and seed yield and improves seed quality of cotton grown on calcareous soils. *Nutr Cycl Agroecosys* 76:19-28.
- Dugger WM (1973) Functional aspects of boron in plants. *Adv Chem Ser* 123:112-129.
- Edwards AH (2014) The semi-micro Kjeldahl method for the determination of nitrogen in coal. *J Appl Chem* 4(6):330-340.
- El-Geddawy Dalia IH, Makhlof BSI (2015) Effect of hill spacing and nitrogen and boron fertilization levels on yield and quality attributes in sugar beet. *Minufiya J Agric Res* 4(1):959-980.
- El-Geddawy IH, Abd-Elhakim AM, Saif LM (2000) Multivariate analysis of yield and relative contribution of variables to its variation under some cultural practices in sugar beet. *Egypt J Basic Appl Sci* 78:449-469.
- Abd El-hady M (2017) Response of sugar beet growth, productivity and quality to foliar application of different forms of boron microelement and number of sprays under new reclaimed soil conditions. *Egypt J Basic Appl Sci* 39(3):401-410.
- El-Sherbeny SE, Khalil M, Hussepn MS (2007) Growth and productivity of rue (*Ruts graveolens*) under different foliar fertilizers application. *J Appl Sci Res* 3(5):399-407.
- Gemici M, Aktas LY, Turkyilmaz B, Guven A (2003) The effects of the boron application on indole-3-acetic acid levels in *Triticum durum* Desf. cv. Gediz seedlings. *CÜ Fen-Edebiyat Fakültesi* 23(2):17-24.
- Gezgin S, Hamurcu M, Apaydin M (2001). Effect of boron application on the yield and quality of sugar beet. *Turk J Agric For* 25:89-95.
- Grzegorzewski K, Ciećko Z, Szostek R (2017) Influence of mineral fertilization on the yield and macroelement content in sugar beet. *Acta Agrophys* 24(2):221-237.
- Hellal FA, Taalab AS, Safaa AM (2009) Influence of nitrogen and boron nutrition on nutrient balance and sugar beet yield grown in calcareous soil. *Ozean J Appl Sci* 2(1):1-10.
- Hirparan DV, Sakarvadia HL, Savaliya CM, Ranpariya VS, Modhavadiya VL (2018) Effect of different levels of boron and molybdenum on growth and yield of summer groundnut (*Arachis hypogaea* L.) under medium black calcareous soils of south Saurashtra region of Gujarat. *Int J Chem Stud* 5(5):1290-1293.
- Iqbal S, Farooq M, Cheema SA, Afzal I (2017) Boron seed priming improves the seedling emergence, growth, grain yield and grain biofortification of bread wheat. *Int J Agric Biol* 19:177-182.
- Jifon JL, Syvertsen JP, Whaley E (2019) Growth environment and leaf anatomy affect nondestructive estimates of chlorophyll and nitrogen in citrus sp. leaves. *J Am Soc Hortic Sci* 130(2):152-158.
- Kristek A, Stojic B, Kristek S (2006) Effect of the foliar boron fertilization on sugar beet root yield and quality. *Poljopriverda* 12(1):22-26.
- Le-Docte AM (1927) Commercial determination of sugar in the beet root using the sacksle-docte process. *Int Sugar J* 29:488-492.
- Makhlof BSI, Abd El-All AEA (2017) Effect of deficit irrigation, nitrogen and potassium fertilization on sugar beet productivity in sandy soils. *Menoufia J Plant Prod* 2:325-346.
- Mirschel W (2018) Manual Leaf Area Measurement on Individual Sugar Beet Plants Taking Plant Density and Irrigation into Account. Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany.
- Mohammadian R, Ghasemi H, Bazrafshan M, Moharramzadah M (2014) Identification of morpho-physiological traits affecting white sugar yield in sugar beet. *J Plant Physiol Breed* 4(2):23-34.
- Nadeem F, Farooq M, Nawaz A, Ahmad R (2019) Boron improves productivity and profit ability of bread wheat

- under zero and plough tillage on alkaline calcareous soil. *Field Crops Res* 239:1-9.
- Nalini P, Gupta B (2013) The impact of foliar boron sprays on reproductive biology and seed quality of black gram. *J Trace Elem Med Biol* 27:58-64.
- Nyomora AMS, Brown PH, Pinney K, Polito VS (2019) Foliar application of boron to almond trees affects pollen quality. *J Am Soc Hortic Sci* 125(2):265-270.
- Perica S, Brown PH, Connell JH, Nyomora AMS, Dordas C, Hu H, Stangoulis J (2001) Foliar boron application improves flower fertility and fruit set of olive. *Sci Hortic* 36(4):714-716.
- Prasad TNVKV, Sudhakar P, Sreenivasulu Y, Latha P, Munaswamy V, Reddy, KR, Sreeprasad TS, Sajanlal PR, Pradeep T (2012) Effect of nanoscale zinc oxide particles on the germination, growth and yield of peanut. *J Plant Nutr* 35(6):905-927.
- Rawashdeh H, Sala F (2013) Effect of different levels of boron and iron foliar application on growth parameters of wheat seedlings. *Afric Crop Sci Conf Proceed* 11:861-864.
- Rehman A, Farooq M, Rashid A, Nadeem F, Stuerz S, Asch F, Bell RW, Siddique KHM (2018) Boron nutrition of rice in different production systems. A review. *Agron Sustain Dev* 38:25.
- Saadati S, Moallemi N, Mortazavi SMH, Seyyednejad SM (2013) Effects of zinc and boron foliar application on soluble carbohydrate and oil contents of three olive cultivars during fruit ripening. *Sci Hortic* 164:30-34.
- Singh Z, Ghosh G, Debbarma V (2017) Effect of different levels of nitrogen, sulphur and foliar application of boron in sunflower (*Helianthus annuus* L.). *Int J Curr Microbiol Appl Sci* 6(10):1336-1342.
- Tambussi EA, Nogués S, Araus JL (2005) Ear of durum wheat under water stress: water relations and photosynthetic metabolism. *Planta* 221(3):446-458.
- Tlili A, Dridi I, Fatnassi S, Hamrouni H, Gueddari M (2019) Boron characterization, distribution in particle-size fractions, and its adsorption-desorption process in a semiarid Tunisian soil. *J Chem Article ID* 2508489, 1-8.
- Ullah S, Khan AS, Malik AU, Afzal I, Shahid M, Razzaq K (2013) Foliar application of boron influences the leaf mineral status, vegetative and reproductive growth, yield and fruit quality of 'Kinnow' mandarin. *J Plant Nutr* 36(10):1479-1495.
- Wanas HA (2002) Petrography, geochemistry and primary origin of spheroidal dolomite from the upper Cretaceous/Lower Tertiary Maghra El-Bahari Formation at Gabal Ataqa, Northwest Gulf of Suez, Egypt. *Sediment Geol* 151(3-4):211-224.

